

## Pediatric Allergy and Clinical Immunology Consult and referral guidelines

## Introduction

We care for children and teens from birth to 18 years. The most common reasons patients are referred include:

- Recurrent Cough or Wheeze
- Recurrent Bronchiolitis or Bronchitis
- Asthma
- Food Allergy
- Immunodeficiency/concerns for frequent infections
- Allergic Rhinitis/Chronic Rhinitis
- Atopic Dermatitis/Eczema
- Urticarial/Angioedema
- Anaphylaxis
- Drug Allergy
- Hypereosinophilia
- Venom Allergy

We want to make referrals easy, fast and efficient for primary care providers. This tool was developed to help create productive visits for you and your patient.

Each guideline includes three sections: suggested work-up and initial management, when to refer, and information needed. Suggested work-ups may not apply to all patients, but these are studies we generally consider during office visits.

Special note: We prefer to look at all imaging studies at each visit. If radiology services were obtained outside of Spectrum Health, we ask the patient's family to provide images on a CD.

Feedback regarding these guidelines is encouraged. Please contact HDVCH Direct to share feedback.

For access to all pediatric guidelines, visit helendevoschildrens.org and type "guidelines" in the search field.



## Appointment priority guide

Immediate	Call HDVCH Direct and/or send to the closest emergency department. Contact HDVCH Direct at (616) 391-2345
	and ask to speak to the on-call provider.
Urgent	Likely to receive an appointment within 48 hours. Call HDVCH Direct and ask to speak to the on-call provider.
Routine	Will receive first available appointment. Send referral via Epic Care Link, fax completed referral form to (616) 267-
	2401, or send referral through Great Lakes Health Connect.

Diagnosis/symptoms	Suggested work-up/initial management	When to refer	Information needed
Recurrent Cough or Wheeze Recurrent Bronchiolitis or Bronchitis	Chest x-ray: PA and lateral Sweat chloride at an accredited CF Center  Trial of bronchodilators at any age If bronchodilators non- responsive, trial of oral and/ or inhaled corticosteroids or Singular® (if age appropriate)  Oral prednisone is typically dosed ~2 mg/kg/day x 5 days minimum	Has been hospitalized, intubated, or frequent ER visits  Frequent need for oral steroid bursts  Less than 2 years old  Unresponsive to usual therapy with increasing medication use  Complicating conditions such as rhinitis, sinusitis, GE-reflux, and/ or pneumonia  Abnormal spirometry or needs frequent monitoring with spirometry	Request for consult and chief concern  Summary of previous treatments and response  Respiratory history since birth, lab results, and chest films  If sweat chloride test was obtained, must be from CF Center accredited lab.
Asthma	None	Same as above	Same as above



Diagnosis/symptoms	Suggested work-up/initial management	When to refer	Information needed
Food Allergy	No testing needed prior to visit	Any question of food allergy	Summary of all previous reactions
	Prescribe/instruct on use of epinephrine auto-injector	History of anaphylaxis	Summary from an ER visits
	Of Note: We do not recommend IgE food allergy broad-range panels. These have a high false positive rate and can lead to false	We recommend all patients with food allergy have care established with an Allergist	Summary of any previous allergy testing
	diagnosis and potential harm to the patient. If testing is pursued, specific IgE to single food groups based on history is preferred. IgG to food has been shown to be of no clinical value in food allergy.	Urgent referral for patient <1 year of age and history of severe eczema/food allergy, as literature shows we may have the opportunity to prevent food allergy in these patients.	



Diagnosis/symptoms	Suggested work-up/initial management	When to refer	Information needed
Immunodeficiency/Concern for Frequent Infections	CBC with diff, IgG, IgA, IgM, HIV	Urgent referral available for concern of serious immunodeficiency	Brief summary of infections and hospitalizations
		· · · · · · · · · · · · · · · · · · ·	All previous radiology results (including CD of film if not done
		Two or more serious sinus infections in 1 year	at a Spectrum Health facility)
		Two or more months on antibiotics with little effect	All culture and lab results outside of Spectrum Health
		Two or more CXR proven pneumonias in 1 year	
		Failure of an infant to gain weight or grow normally	
		Recurrent, deep skin or organ abscesses	
		Persistent thrush in mouth or fungal infection of skin	
		Need for IV antibiotics to clear infections	
		Two or more deep seated infections, including septicemia	
		Family history of primary immunodeficiency Infection with rare or low virulent organisms	
		Unexplained bronchiectasis	



management		
Trial of second generation H-1 antihistamines (Zyrtec/cetirizine or Allegra/fexofenadine) at any age	Symptoms refractory to antihistamine and nasal steroid	History of symptoms and therapies to this point
Trial nasal steroid if tolerated	Need to clarify diagnosis of allergy vs non-allergic	
	Need to identify specific allergens for environmental management Need for evaluation for allergy shots	
Topical corticosteroids (cream/ointment, not lotion)  Frequent emollients  Oral H-1 antihistamine at night	Continued flares of atop dermatitis despite current treatment  Urgent referral for all patients <1 year of age with severe eczema to evaluate for early introduction/ prevention of food allergy in accordance with LEAP study, EAT study, and 2016 food allergy practice parameters.	History of previous treatment  Any labs that were obtained outside of Spectrum Health
- (	Allegra/fexofenadine) at any age  Trial nasal steroid if tolerated  Topical corticosteroids (cream/ pintment, not lotion)  Frequent emollients	Allegra/fexofenadine) at any age  Trial nasal steroid if tolerated  Need to clarify diagnosis of allergy vs non-allergic  Need to identify specific allergens for environmental management Need for evaluation for allergy shots  Topical corticosteroids (cream/pointment, not lotion)  Continued flares of atop dermatitis despite current treatment  Frequent emollients  Urgent referral for all patients <1 year of age with severe eczema to evaluate for early introduction/prevention of food allergy in accordance with LEAP study, EAT study, and 2016 food allergy



Diagnosis/symptoms	Suggested work-up/initial management	When to refer	Information needed
Diagnosis/symptoms Urticaria/Angioedema		When to refer  Unexplained acute urticaria  Symptoms that are refractory, and continue despite BID H-1 and H-2 antihistamine  Angioedema, without urticarial, accompanied by low C4 (concern for Hereditary Angioedema)	History of previous treatment  Any labs that were obtained outside of Spectrum Health
	Angioedema/swelling without urticaria		



Diagnosis/symptoms	Suggested work-up/initial management	When to refer	Information needed
Anaphylaxis	Could consider baseline tryptase	Any case of anaphylaxis, especially unexplained, should be referred to an allergist	Brief history of anaphylactic event  Any labs obtained, especially tryptase, if this is obtained during ER visit for anaphylaxis
Drug Allergy	Due to high rate of false negatives - unless needed for urgent/emergent reasons (i.e. chemotherapy) - we cannot test to drugs until 6 weeks after reaction	History of allergy/reaction to a medication that is medically indicated for the patient to take in the future  History of allergy/reaction to multiple medications that make prescribing future medications difficult  Any history of penicillin allergy in children >10 years old	History of reaction to all medications
Hypereosinophilia	CBC with diff  Stool o/p, toxocara canis antibody, and Strongyloides (note, there is risk of death if prednisone is given to patient with strongyloides)	Absolute eosinophil count >1000, with negative stool o/p, toxocara canis antibody, and strongyloides antibody	All laboratory results outside of Spectrum Health (including all CBCs that have been obtained)
Venom Allergy	Prescribe injectable epinephrine	All patients with history of reaction to stinging insect that is more than a large reaction at the site of the sting/bite.	Brief history of reaction

## **HDVCH Direct phone: (616) 391-2345**

HDVCH developed these referral guidelines as a general reference to assist referring providers. Pediatric medical needs are complex and these guidelines may not apply in every case. HDVCH relies on its referring providers to exercise their own professional judgment with regard to the appropriate treatment and management of their patients. Referring providers are solely responsible for confirming accuracy, timeliness, completeness, appropriateness and helpfulness of this material and making all medical, diagnostic and prescription decisions.